

## Enfamil® ProSobee® Infant Formula Powder Dilutions Per 1 Scoop

**Note: Powdered products are not commercially sterile and should not be used for immunocompromised patients, unless clinically required, and then under strict medical supervision of preparation and use.**

Calories Desired, per fl oz	1 Scoop Powder added to indicated Volume of Water*, fl oz	Formula Yield, fl oz
20	2.0	2.2
22	1.8	2.0
24	1.6	1.9
26	1.5	1.7
27	1.4	1.7
28	1.4	1.6
30	1.3	1.5

Enfamil ProSobee Powder should be measured with **unpacked**, level scoops, using the scoop enclosed in the can.

\*Fluid ounce measures are provided for convenience when larger variance in prepared dilution is acceptable. Fluid ounce measures in above table are rounded to nearest 0.1 fluid ounce.

One-half fl oz of water can be measured using one tablespoon (1/2 fl oz water = 1 Tbsp water).

When accuracy is essential, water should be measured in mL, and powder should be weighed rather than measured. Contact [mjmedicalaffairs@reckitt.com](mailto:mjmedicalaffairs@reckitt.com) for those instructions.

Powders mix best when added on top of water. Then close bottle and shake for about 5 seconds. The number of scoops and the amount of water to add can be doubled or tripled to make a larger volume.

One scoop of Enfamil ProSobee Powder (8.8 g) provides 45 Calories.

One scoop of Enfamil ProSobee Powder displaces about 0.2 fl oz of water.

One fluid ounce = 29.57 mL.