

PurAmino™ Infant Formula

Preparation at Various Concentrations Using 1 Scoop of Powder

Note: Powdered products are not commercially sterile and should not be used for premature or immunocompromised patients, unless clinically required, and then under strict medical supervision of preparation and use.

Calories Desired per fluid ounce	Volume of Water** -Add 1 unpacked level scoop (4.5 g) of powder to indicated volume of water		Formula Yield	
	mL	fl oz	mL	fl oz
20 Cal/fl oz	30	1	33	1.1
22 Cal/fl oz	27	0.9	30	1
24 Cal/fl oz	24	0.8	28	0.9
26 Cal/fl oz	22	0.7	26	0.9
27 Cal/fl oz	21	0.7	25	0.8
28 Cal/fl oz	20	0.7	24	0.8
30 Cal/fl oz	19	0.6	22	0.8

**When accuracy is essential, mL measurements should be used for water. Fluid ounce measures are provided for convenience when a larger variance in prepared concentration is acceptable. Fluid ounce measures in above table are rounded to the nearest 0.1 fl oz.

- One fl oz = 29.57 mL.
- ½ fl oz of water can be measured using one tablespoon (1 Tbsp water = ½ fl oz water)
- Powders mix best when added on top of water.
- The number of scoops and the amount of water to add can be doubled or tripled to make a larger volume.
- One unpacked level scoop of PurAmino™ infant formula powder (4.5 g) provides 23 calories.
- Each gram of PurAmino™ infant formula powder displaces 0.77 mL water.