Drug Facts

Active ingredients (in each lozenge) — Purposes

Dextromethorphan HBr 5.0 mg ..... Cough suppressant
Hexylresorcinol 2.0 mg ..... Oral pain reliever

Uses temporarily relieves

- cough due to minor throat and bronchial irritation as may occur with the common cold
- occasional irritation and pain due to a sore mouth or sore throat
- pain associated with canker sores

Warnings

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- in children under 6 years of age

When using this product do not exceed recommended dosage

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor or dentist if

- sore mouth symptoms do not improve in 7 days
- irritation, pain, or redness persists or worsens
- swelling develops
- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions dose as follows or as directed by a doctor or dentist

adults and children 12 years of age and older	take 2 lozenges (one immediately after the other) and allow each lozenge to dissolve slowly in the mouth; may be repeated every 4 hours as needed, not to exceed 12 lozenges in any 24-hour period
children 6 to under 12 years of age	take 1 lozenge and allow product to dissolve slowly in the mouth; may be repeated every 4 hours as needed, not to exceed 6 lozenges in any 24-hour period
children under 6 years of age	do not use

Other information

- store at 20-25°C (68-77°F)
- protect contents from humidity

Inactive ingredients corn syrup, D&C yellow no. 10, ethyl lactate, FD&C blue no. 1, flavor, propylene glycol, sucrose, water

Questions? 1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.

Mucinex®